

Assisted tunneling of a wave packet between square barriers

G. Kälbermann*

*Soil and Water department, The Robert H. Smith Faculty of Agriculture,
Food and Environment Hebrew University, Rehovot 76100, Israel*

The assisted tunneling of a wave packet between square one dimensional barriers is treated analytically. The survival probability is calculated exactly for a potential mimicking a constant electric field with arbitrary time dependence. The pole spectrum of the properly normalized unperturbed wavefunctions determines the decay time. The tunneling probability is enhanced by the perturbation. The results are exemplified for a simplified model of nuclear α decay.

PACS numbers: 03.65.Xp, 73.43.Jn 23.60.+e

I. INTRODUCTION

The quantum tunneling paradigm explains diverse processes, as currents in Josephson junctions and nuclear α decay. Energetically forbidden regions for a classical object, become accessible for a quantum system.

Gurney and Condon and Gamow[2, 3], used the tunneling model to reproduce the experimentally measured huge range of α decay lifetimes of similar unstable nuclei. The scheme has withstood continuous scrutiny in the intervening years [4, 7].

In a previous work we investigated the tunneling of a one dimensional metastable state between delta function barriers, excited by a time dependent potential.[1] By inspecting the modification of the spectrum of the poles ruling the decay process, we estimated a non-negligible enhancement of the tunneling probability when an external time dependent perturbation acts on the system.

In the present work we find exact solutions to the assisted tunneling process, with square barrier potentials instead of delta functions. Square potentials provide a more realistic description of the barriers involved in decay processes.

Decay times are enormously different from the natural time scales of nuclear and even atomic phenomena. It is not possible to follow the decay process numerically. Analytical expressions are then of the utmost importance[1, 5, 6]. The method and formulas presented here aim at contributing in this direction. In the next section we present the model and the analytical solution method. In section 3 we solve the equations for a time harmonic, linear potential and evaluate the effect of the perturbation on the tunneling process. Some mathematical details are presented in appendix A.

II. ANALYTICAL METHOD FOR ASSISTED TUNNELING

The Schrödinger equation for a one dimensional system between square barriers is

$$i \frac{\partial \Psi}{\partial t} = \frac{-1}{2m} \frac{\partial^2 \Psi}{\partial x^2} + \gamma (\Theta(d - |x|)\Theta(|x| - x_0))\Psi \quad (1)$$

In the following all our units are either fm for length and time, and fm^{-1} for energies, momenta and potential strength γ .

We consider an initial Gaussian wavepacket located $t = 0$ in the region between the barriers.

$$\Psi(x, t = 0) = e^{-\frac{x^2}{\Delta^2}} \quad (2)$$

where Δ is the width parameter of the packet. This packet is a reasonable model for the metastable state and facilitates the analytical evaluation of the decay amplitudes. However, the method presented here is not limited to gaussian packets.

Inside the barrier region the packet disperses if $\Delta < x_0$, but, soon enough swelling comes to a halt due to the presence of the barriers. It can then spread only through the tunneling process governed by the barriers.

The even $\chi_e(x) = \frac{\varphi_e(k)}{\sqrt{\pi}n_e(k)}$ and odd $\chi_o(x) = \frac{\varphi_o(k)}{\sqrt{\pi}n_o(k)}$ stationary states of eq.(1) for energies below the barrier height γ are

$$\varphi_e(k) = \begin{cases} \cos(kx); & |x| < x_0 \\ A_1 e^{\kappa|x|} + B_1 e^{-\kappa|x|}; & d > |x| > x_0 \\ C_1 \cos(kx) + \text{sign}(x)D_1 \sin(kx); & |x| > d \end{cases} \quad (3)$$

where κ is defined in eq.(7), and $\text{sign}(x) = 1, -1$ for $x > 0, x < 0$.

$$\varphi_o(k) = \begin{cases} \sin(kx); & |x| < x_0 \\ \text{sign}(x)(A_2 e^{\kappa|x|} + B_2 e^{-\kappa|x|}); & d > |x| > x_0 \\ \text{sign}(x)C_2 \cos(kx) + D_2 \sin(kx); & |x| > d \end{cases} \quad (4)$$

*Electronic address: hope@vms.huji.ac.il

where we have extracted a factor of $\sqrt{\pi}$ from the normalizations for convenience.

The set of even-odd functions is orthonormal and complete.[8, 9] The zeros of the normalization factors govern the exponential decay of the metastable wave functions in the inner region[1].

The normalization factors are given by

$$\begin{aligned}
((n_e(k))^2 &= (C_1(k)^2 + D_1(k)^2) \\
D_1 &= -\frac{1}{2} \left(c_2 \kappa e_2 k s_1 \right. \\
&\quad - c_2 \kappa^2 e_2 c_1 + c_2 \kappa^2 e_1 c_1 \\
&\quad + c_2 \kappa e_1 k s_1 + e_2 k^2 s_1 s_2 - e_2 \kappa c_1 k s_2 \\
&\quad \left. - e_1 \kappa c_1 k s_2 - e_1 k^2 s_1 s_2 \right) / (\kappa k e_3) \\
C_1 &= \frac{1}{2} \left(e_2 k s_1 \kappa s_2 - e_2 \right. \\
&\quad \kappa^2 c_1 s_2 + e_1 \kappa^2 c_1 s_2 + e_1 k s_1 \kappa s_2 \\
&\quad - c_2 e_2 k^2 s_1 + c_2 e_2 \kappa c_1 k + c_2 e_1 \kappa c_1 k \\
&\quad \left. + c_2 e_1 k^2 s_1 \right) / (\kappa k e_3) \quad (5)
\end{aligned}$$

$$\begin{aligned}
(n_o(k))^2 &= (C_2(k)^2 + D_2(k)^2) \\
D_2 &= -\frac{1}{2} \left(-c_2 e_2 \kappa c_1 k \right. \\
&\quad - c_2 \kappa^2 e_2 s_1 + c_2 \kappa^2 e_1 s_1 \\
&\quad - c_2 e_1 \kappa c_1 k - e_2 k^2 c_1 s_2 - e_2 k s_1 \kappa s_2 \\
&\quad \left. - e_1 k s_1 \kappa s_2 + e_1 k^2 c_1 s_2 \right) / (\kappa k e_3) \\
C_2 &= -\frac{1}{2} \left(e_2 \kappa c_1 k s_2 \right. \\
&\quad + e_2 \kappa^2 s_1 s_2 - e_1 \kappa^2 s_1 s_2 \\
&\quad + e_1 \kappa c_1 k s_2 - c_2 e_2 k^2 c_1 - c_2 \kappa e_2 k s_1 \\
&\quad \left. - c_2 \kappa e_1 k s_1 + c_2 e_1 k^2 c_1 \right) / \\
&\quad (\kappa k e_3) \quad (6)
\end{aligned}$$

where

$$\begin{aligned}
\kappa &= \sqrt{2 m \gamma - k^2} & e_3 &= e^{\kappa (x_0 + d)} \\
e_2 &= e^{2 \kappa d} & e_1 &= e^{2 \kappa x_0} \\
c_2 &= \cos(k d) & s_2 &= \sin(k d) \\
c_1 &= \cos(k x_0) & s_1 &= \sin(k x_0)
\end{aligned} \quad (7)$$

We assume a time dependent spatially linear perturbation potential

$$V(x, t) = \mu x G(t) \quad (8)$$

with μ a coupling constant. For a spatially constant time-harmonic electric field of intensity E_0 interacting with an α particle of charge $2|e|$, we have

$$\mu G(t) = 2|e| E_0 \sin(\omega t) \quad (9)$$

In order to eliminate the explicit space dependence of the perturbation term in the Schrödinger equation, we apply the unitary transformation

$$\begin{aligned}
\Psi(x, t) &= e^{-i\sigma} \Phi \\
\sigma &= \mu x \zeta + \int \frac{\zeta^2 \mu^2}{2 m} dt \\
\zeta &= \int G(t) dt \quad (10)
\end{aligned}$$

The Schrödinger equation(1) including the perturbation of eq.(8) becomes

$$\begin{aligned}
i \frac{\partial \Phi}{\partial t} &= \frac{-1}{2 m} \frac{\partial^2 \Phi}{\partial x^2} + \gamma (\Theta(d - |x|) \Theta(|x| - x_0)) \Phi \\
&\quad + i \zeta(t) \frac{\mu}{m} \frac{\partial \Phi}{\partial x} \quad (11)
\end{aligned}$$

The wave function Φ is expanded in the complete set of even and odd states of the unperturbed Schrödinger equation (1) including states with energies above the barrier

$$\Phi(x, t) = \sum_{i=e,o} \int_0^\infty \chi_i(k, x) c_i(k, t) e^{\frac{-i k^2}{2 m} t} dk \quad (12)$$

The amplitudes $c_{e,o}$ consequently obey the time evolution

$$\begin{aligned}
\dot{c}_o(k) &= -\frac{\mu k \zeta}{m n_e(k) n_o(k)} c_e(k) \\
\dot{c}_e(k) &= \frac{\mu k \zeta}{m n_e(k) n_o(k)} c_o(k) \quad (13)
\end{aligned}$$

where we have used the superposition integral calculated in appendix A [12]

$$\begin{aligned}
I &= \int e^{-\frac{i(k'^2 - k^2)t}{2m}} \chi_e(k, x) \frac{\partial \chi_o(k', x)}{\partial x} dx \\
&= \frac{k}{n_e(k) n_o(k)} \delta(k - k') \quad (14)
\end{aligned}$$

and a dot denotes a time derivative.

The exact solution to eqs.(13) for an initially symmetric wave packet such as the one of eq.(2) is

$$\begin{aligned}
c_e(k, t) &= c_e(k, 0) \cos\left(\frac{\mu k Y}{m n_e n_o}\right) \\
c_o(k, t) &= -c_e(k, 0) \sin\left(\frac{\mu k Y}{m n_e n_o}\right) \\
Y(t) &= \int_0^t \zeta(t') dt' \\
c_e(k, 0) &= \frac{\Delta}{\pi^{\frac{1}{4}} n_e(k)} e^{-\frac{k^2}{4 \Delta^2}}
\end{aligned} \tag{15}$$

where $c_e(k, 0)$ corresponds a well localized $\Delta \ll x_0$ wave packet of the form of eq.(2). Other initial wave functions merely change the functional dependence of $c_e(k, 0)$.

III. ASSISTED TUNNELING SOLUTION

We are interested in the decay of the wave packet from the internal zone leaking out to infinity. As discussed in [1], the most important contribution to the integral in eq.(12) originates from the poles in the amplitudes of eq.(15). This is indeed correct for the unperturbed decay process. However, for the perturbed process, the amplitudes are oscillatory. The poles lie in extremely close proximity to the real momentum axis. When the integration over this variable is effected, the argument of the harmonic functions in eq.(15) can oscillate wildly, suppressing the contribution of the poles.(see figures 2 and 3 below) However, if the argument of the harmonic functions is of order one, the poles will still dominate. This can be achieved by choosing a sufficiently high frequency for the external perturbation of eq.(9).

The normalization factors around their minima can be cast in the form

$$k^2 n_{e,o}^2 \approx \lambda_{j,(e,o)} (k^2 - k_{j,(e,o)}^2)^2 + \beta_{j,(e,o)} \tag{16}$$

where j enumerates the zeros of n_e, n_o .

The minima are located close to

$$\begin{aligned}
k_e &\approx \frac{(2n+1) \pi p}{2 (1+p x_0)} \\
k_o &\approx \frac{n \pi p}{(1+p x_0)} \\
p &= \sqrt{2 m \gamma}
\end{aligned} \tag{17}$$

Eqs.(17) provide starting values for the numerical search of the location of the poles of the wave function. The poles in the complex momentum plane, appear in pairs located symmetrically above and below the real momentum axis. The stationary wave functions of eqs.(3,4) consist of incoming and outgoing parts. The standard S matrix, reflecting the behavior of the outgoing boundary condition at spatial infinity, has poles in the lower half of




FIG. 1: $10^{-10} n_e(k)^{-1}$, and $10^{-10} n_o(k)^{-1}$, as a function of k in units of fm^{-1} for a typical α decay barrier

the complex momentum plane, whereas its inverse, corresponding to the incoming condition, has poles in the upper half. Consequently, both sets of poles in the upper and lower complex momentum plane appear when incoming and outgoing pieces are present.[10].

As the poles are extremely sharp in position, it was necessary to use a quadruple precision algorithm on an α cluster mainframe to find them. The imaginary parts of the poles are typically many orders of magnitude smaller than the real parts.

Figure 1 shows $10^{-10} n_e(k)^{-1}$, and $10^{-10} n_o(k)^{-1}$, for the parameters $m = 3727 MeV = 18.88 fm^{-1}$, $x_0 = 12 fm$, $d = 22 fm$, $\gamma = 22 MeV = 0.11 fm^{-1}$, corresponding to a typical α decay barrier. The spikes are due to the extreme closeness of the minima of the normalization factors to their complex zeros.

When eq.(12) is evaluated by contour integration in the complex k^2 plane, the contour has to be closed from below. In the lower half-plane the convergence is insured by the exponential $e^{-i \frac{k^2 t}{2m}}$. The poles are separated from each other and the integrand drops essentially to zero outside the pole, hence, the wave function consists of an incoherent sum of the residues of the different poles. The negative imaginary part of each pole induces a time decaying exponential. Each exponent determines a different decay constant and decay time.[1] If the original wave function is even in space, as in eq.(2), only even poles show up in the unperturbed wave function.

The argument of the harmonic functions in eq.(15) includes a potentially large denominator near the minima of the normalization factors. The product of these even-odd factors near a minimum is depicted in figures 2 and 3. This product is not only finite, but of order one. We were not able to produce an analytical formula for this product, although the numerical values are very suggestive.




FIG. 2: $n_o(k_{j,e}) n_{j,e}^{-1}$, around the minima of $n_e(k)$. Parameters as in figure 1

Figure 2 depicts $(n_e(k) n_o(k))^{-1}$, around the minima of $n_e(k)$. Note the degree of accuracy needed for the location of the minima. The parameters are those of figure 1. Figure 3 depicts $(n_e(k) n_o(k))^{-1}$ around the minima of $n_o(k)$.

Expanding the harmonic functions in eq.(15), the exact expression for the wave function in the region between the barriers becomes

$$\Phi(x, t) = \Phi_{ee}(x, t) + \Phi_{oo}(x, t) + \Phi_{eo}(x, t) + \Phi_{oe}(x, t) \quad (18)$$

where

$$\begin{aligned} \Phi_{ee}(x, t) &= \sum_j \sum_{n=0}^{\infty} G_j(x, t) \frac{(u_{j,e})^n}{(n!)^2} \\ G_j(x, t) &= \frac{\pi^{3/4} \Delta \cos(s_{j,e} x) s_{j,e} e^w}{2 \sqrt{\beta_{j,e} \lambda_{j,e}}} \end{aligned}$$




FIG. 3: $n_o(k_{j,e}) n_{j,e}^{-1}$, around the minima of $n_o(k)$. Parameters as in figure 1

$$\begin{aligned} u_{j,e} &= -\left(\frac{\mu q_{j,e} Y(t)}{2 m n_o(q_{j,e}) \sqrt{\beta_{j,e}}} \right)^2 \\ w &= -q_{j,e} (\Delta^2/4 + i t/(2 m)) \\ q_{j,e} &= (k_{j,e})^2 - i \frac{\beta_{j,e}}{\lambda_{j,e}} \\ s_{j,e} &= \sqrt{q_{j,e}} \end{aligned} \quad (19)$$

$$\begin{aligned} \Phi_{oo}(x, t) &= \sum_j \sum_{n=0}^{\infty} H_j(x, t) \frac{(u_{j,o})^n}{(n!)^2 (2n+1)} \\ H_j(x, t) &= -\frac{\pi^{3/4} \Delta \sin(s_{j,o} x) \mu q_{j,o} Y(t) e^w}{2 m n_e(q_{j,o})^2 \sqrt{\beta_{j,o} \lambda_{j,o}}} \\ u_{j,o} &= -\left(\frac{\mu q_{j,o} Y(t)}{2 m n_e(q_{j,o}) \sqrt{\beta_{j,o}}} \right)^2 \\ q_{j,o} &= (k_{j,o})^2 - i \frac{\beta_{j,o}}{\lambda_{j,o}} \\ s_{j,o} &= \sqrt{q_{j,o}} \end{aligned} \quad (20)$$

$$\Phi_{eo}(x, t) = \sum_j \sum_{n=1}^{\infty} L_j(x, t) \frac{(u_{j,o})^n}{((n-1)!)^2 n (2n-1)}$$

$$L_j(x, t) = \frac{2\pi^{3/4} \Delta \cos(s_{j,o} x) \sqrt{\beta_{j,o}} e^w}{2 m n_e (q_{j,o})^2 s_{j,o} \lambda_{j,o}} \quad (21)$$

$$\Phi_{oe}(x, t) = \sum_j \sum_{n=0}^{\infty} M_j(x, t) \frac{(u_{j,e})^n}{(2n+1)(n!)^2}$$

$$M_j(x, t) = \frac{\pi^{3/4} \mu Y(t) s_{j,e} \Delta \cos(s_{j,e} x) e^w}{2 m n_o (q_{j,e})^2 \sqrt{\beta_{j,e}} \lambda_{j,e}} \quad (22)$$

The series in eqs.(19,20,21,22) can be expressed in terms of standard functions

$$\Phi_{ee}(x, t) = \sum_j G_j(x, t) J_0(u_{j,e})$$

$$\Phi_{oo}(x, t) = \sum_j H_j(x, t) \left[J_0(u_{j,o}) - \frac{\pi}{2} \left(J_0(u_{j,o}) \mathbf{H}_1(u_{j,o}) - J_1(u_{j,o}) \mathbf{H}_0(u_{j,o}) \right) \right]$$

$$\Phi_{eo}(x, t) = \sum_j L_j(x, t) \left[J_0(u_{j,o}) - \frac{\pi}{2} \left(J_0(u_{j,o}) \mathbf{H}_1(u_{j,o}) - J_1(u_{j,o}) \mathbf{H}_0(u_{j,o}) \right) \right]$$

$$\Phi_{oe}(x, t) = \sum_j M_j(x, t) \left[2 J_0(u_{j,e}) - \frac{2 J_1(u_{j,e})}{u_{j,e}} - \pi \left(J_0(u_{j,e}) \mathbf{H}_1(u_{j,e}) - J_1(u_{j,e}) \mathbf{H}_0(u_{j,e}) \right) \right] \quad (23)$$

where $J_{0,1}$ denotes the Bessel function of the order (0,1), and $\mathbf{H}_{0,1}$ is the Struve $\mathbf{H}$ function. [11]

For long times we can use the asymptotic expansions of the Bessel and Struve functions to obtain[11]

$$\Phi_{ee}(x, t) \rightarrow \sum_j G_j(x, t) \left(\sqrt{\frac{2}{u_{j,e} \pi}} \sin(u_{j,e} + \frac{\pi}{4}) + O(u_{j,e}^{-\frac{3}{2}}) \right)$$

$$\Phi_{oo}(x, t) \rightarrow \sum_j H_j(x, t) \left(\frac{1}{u_{j,o}} + O(u_{j,o}^{-\frac{3}{2}}) \right)$$

$$\Phi_{eo}(x, t) \rightarrow \sum_j L_j(x, t) \left(\frac{1}{u_{j,o}} + O(u_{j,o}^{-\frac{3}{2}}) \right)$$

$$\Phi_{oe}(x, t) \rightarrow \sum_j M_j(x, t) \left(\frac{2}{u_{j,e}} + O(u_{j,e}^{-\frac{3}{2}}) \right) \quad (24)$$

From eq.(24) it is clear that for long times the even-even component is essentially the dominant one. We can now




FIG. 4: $S(t)$ of eq.(25), without, full line, and with harmonic perturbation, dashed line, see text for parameters

evaluate the influence of the perturbation on the decay time. Figure 4 depicts the nonescape probability of the system in the region between the barriers

$$S(t) = \int_{-x_0}^{x_0} |\Psi(x, t)|^2 dx \quad (25)$$

for the square barrier case, we have used the parameters relevant for α decay mentioned below eq.(1) and an initial wave packet of width $\Delta = 4fm$. The lower curve corresponds to (see eq.(9)) an electric field of $E = 0.1V/m$, and a frequency of $\omega = 10\pi \cdot 10^{17}Hz$. The aim of this work is to present the analytical solutions and show the acceleration of the decay process. A more realistic choice of parameters demands a thorough numerical evaluation of the integrals outside the poles and is postponed for future work. For a time harmonic perturbation(9), there are two major corrections to the unperturbed decay rate. The amplitude of the wave function is to lowest order multiplied by a factor of the form $(1 - ct^2/4)$, with $c \approx \frac{\mu^2 k^2}{m^2 \omega^2}$ from the series in $u_{j,e}$. This factor depletes the wave at times of the order of the period of the harmonic perturbation for moderate values of μ , much faster than the actual decay time determined by the poles. A second correction already noted in[1], arises when considering the higher order poles $j > 1$ in the sum of eq.(19). The latter generates a different decay time for each pole[1].




FIG. 5: $S(t)$ of eq.(25), without, full line, and with harmonic perturbation, dashed line, for δ barriers

Figure 5 shows the results for delta function barriers $\nu \delta(x-x_0) + \nu \delta(x+x_0)$, with parameters corresponding to the square barrier case $\nu = \gamma(d-x_0)$. The unperturbed decay time of around 10^{-16} sec is orders of magnitude smaller than the square barrier one. The assisted tunneling is barely noticeable.

In summary, we presented a simple analytical method to calculate assisted tunneling. The technique can be generalized to three dimensions, and to other potentials and initial metastable states. Such endeavor is worth pursuing due to the dramatic impact assisted tunneling may have on nuclear technologies as well as on solid state devices. The question of electron screening for the nuclear decay acceleration must also be addressed to produce experimentally meaningful predictions.

Acknowledgments

It is a pleasure to acknowledge the anonymous referee's remarks and corrections.

-
- [1] G. Kälbermann, Phys. Rev. **C77** 041601(R)(2008).
 - [2] R. W. Gurney and E.U. Condon, Phys. Rev. **33**,127 (1929).
 - [3] G. Gamow, Zeit. Phys. **51**, 204 (1928).
 - [4] E. L. Medeiros et al. Jour. of Phys. **G32**, B23 (2006).
 - [5] Van Dijk W. and Nogami Y., Phys. Rev. **C65**, 024608 (2002).
 - [6] Garcia-Calderon G., Maldonado I. and Villavicencio J., Phys. Rev. **A76**, 012103 (2007).
 - [7] B. Holstein, Am. J. Phys. **64**, 1062 (1996).
 - [8] K. R. Brownstein, Am. J. Phys. **43**, 173 (1975).
 - [9] S. H. Patil, Am. J. Phys. **68**, 712 (2002).
 - [10] A. I. Baz, Ya. B. Zeldovich and A. M. Perelomov, *Scattering, Reactions and Decay in Nonrelativistic Quantum Mechanics*, Israel Program for Scientific Translations, Jerusalem (1969).
 - [11] M. Abramowitz and I. Stegun, *Handbook of mathematical functions*, Dover Publications, NY (1972).
 - [12] In ref.[1] there was a sign misprint in the second of eqs.(11)

IV. APPENDIX A: PERTURBATION MATRIX ELEMENT

The matrix element of eq.(14) does not vanish for transitions between even and odd states solely. The matrix element evidently diverges and we seek to evaluate the divergent part. The most important contribution comes from the outer region of $|x| > d$. In the inner region there

will be a finite contribution that is negligible as compared to the δ function contribution of the outer region. Consequently we can use the outer region expression throughout.

The functions $\chi_{e,o}$ are then replaced for all x by

$$\begin{aligned}
 \sqrt{\pi} n_e(k) \phi_e(k) &= (E e^{i k x} + E^* e^{-i k x}), x > 0 \\
 \sqrt{\pi} n_e(k) \phi_e(k) &= (E e^{-i k x} + E^* e^{i k x}), x < 0 \\
 \sqrt{\pi} n_o(k) \phi_o(k) &= (F e^{i k x} + F^* e^{-i k x}), x > 0 \\
 \sqrt{\pi} n_o(k) \phi_o(k) &= -(F e^{-i k x} + F^* e^{i k x}), x < 0 \\
 E &= \frac{C_1 - i D_1}{2} \\
 F &= \frac{C_2 - i D_2}{2}
 \end{aligned} \tag{26}$$

Taking advantage of the symmetry properties of the wave functions, the matrix element of eq.(14) becomes

$$\begin{aligned}
 M_{k',k} &= \int_{-\infty}^{\infty} e^{-\frac{i(k'^2 - k^2)x}{2m}} \chi_e(k, x) \frac{\partial \chi_o(k', x)}{\partial x} dx \\
 &= \frac{2}{\pi n_e(k) n_o(k')} \sum_{i=1}^4 (I_i(x \rightarrow \infty) - I_i(x = 0)) \\
 I_1(x) &= i k' E \tilde{F} \frac{e^{i(k+k')x}}{i(k+k')} \\
 I_2(x) &= -i k' E \tilde{F}^* \frac{e^{i(k-k')x}}{i(k-k')}
 \end{aligned}$$

$$\begin{aligned}
I_3(x) &= i k' E^* \tilde{F} \frac{e^{i(-k+k')x}}{i(-k+k')} \\
I_4(x) &= -i k' E^* \tilde{F}^* \frac{e^{-i(k+k')x}}{-i(k+k')} \quad (27)
\end{aligned}$$

where the *tilde* denotes a factor that depends on k' instead of k . The all important contribution comes from $k' \rightarrow k$

A lengthy but straightforward calculation then gives the result of eq.(14)

$$M_{k',k} = \frac{k}{n_e(k) n_o(k)} \delta(k - k') \quad (29)$$

$$M_{k',k} \approx 2 i k' (E^* F - E F^*) \frac{\delta(k - k')}{n_e(k) n_o(k')} \quad (28)$$